(19) INDIA

(22) Date of filing of Application :27/12/2023

(43) Publication Date: 12/01/2024

(54) Title of the invention: METHOD FOR OXIDATIVE DEGRADATION OF SULPHONIC ACID DYES FROM WASTEWATER USING FERRIC PERCHLORATE AS CATALYST

(51) International classification :G06T0005000000, C02F0001720000, C08G0061120000, H04N0019610000,

C02F0001320000

(86) International Application No Filing Date (87) International

(87) International
Publication No
(61) Patent of Addition
to Application Number :NA
:NA

Filing Date
(62) Divisional to
Application Number

NA

:NA

Filing Date :NA

(71)Name of Applicant:

1)Andhra University
Address of Applicant :Andhra University, Waltair,

Visakhapatnam-530003, Andhra Pradesh, India. Visakhapatnam --

Name of Applicant: NA Address of Applicant: NA (72)Name of Inventor: 1)M. R. Aruna Kumari

Address of Applicant :Research Scholar, Department of Chemistry, Andhra university, Waltair, Visakhapatnam-530003, Andhra Pradesh, India. Visakhapatnam ------

Andina Fradesh, mdia. Visakhapatha 2)Prof. P. Shyamala

Address of Applicant: Head of the Department Department of Chemistry, Andhra University, Visakhapatnam, Andhra Pradesh, 530003, India. Visakhapatnam ------

3)B. Leela Kumari

Address of Applicant :Research Scholar, Department of Chemistry, Andhra university, Waltair, Visakhapatnam-530003, Andhra Pradesh, India. Visakhapatnam -----

(57) Abstract:

ABSTRACT: Title: Method for Oxidative Degradation of Sulphonic Acid Dyes from Wastewater Using Ferric Perchlorate as Catalyst The present disclosure proposes a method for oxidative degradation of sulphonic acid dyes from wastewater using ferric perchlorate as catalyst. The method comprises a cylindrical reactor (102), an oxidant (104), a catalyst (106), a mechanical stirrer (108) and an analyzing unit (110). The method that improves the oxidation process but also simplifies the overall operational degradation process. The method that offers cost-effectiveness and wide applicability. The method that enhances the degradation of sulphonic dyes with periodate in presence of Ferric perchlorate is above 300 times much faster with high efficiency. The method that enhances the efficiency of the degradation and removal of sulphonic acid from the wastewater. The method that utilizes ferric perchlorate, a water-soluble homogeneous catalyst to achieve high-efficient degradation and removal of sulphonic acid from the wastewater.

No. of Pages: 20 No. of Claims: 7